

Physics 233
Experiment 9

The Compound Pendulum

References

Stephenson, *Mechanics and Properties of Matter*, Wiley, 1960,
Ch. 6, (QA 807 S82).

Introduction

In this experiment we shall see how the period of oscillation of a compound, or physical, pendulum depends on the distance between the point of suspension and the center of mass.

The compound pendulum you will use in this experiment is a one metre long bar of steel which may be supported at different points along its length, as shown in Fig. 1.




Figure 1 A compound pendulum.

If we denote the distance between the point of suspension, O, and the center of mass, by l , the period of this pendulum is:

$$T = 2 \pi \sqrt{\frac{k^2 + l^2}{gl}} \quad (1)$$

where k is the radius of gyration of the bar about an axis passing through the centre of mass. You should derive this expression.

The period of a simple pendulum of length l' is:

$$T = 2\pi \sqrt{\frac{l}{g}} \quad (2)$$

By equating Eqs (1) and (2) and solving for l , we may find the values of l such that the compound pendulum has the same period as that of a simple pendulum of length l' :

$$l = \frac{l' \pm \sqrt{l'^2 - 4k^2}}{2} \quad (3)$$

As you can see, there are two values of l , which we will label l_1 and l_2 , for which the period of the compound pendulum is the same as that of the given simple pendulum.

There is a value of l for which the compound pendulum has a minimum period. The minimum period may be found from Eq. (1) by setting:

$$\frac{dT}{dl} = 0$$

One finds:

$$T_{\min} = 2\pi \sqrt{\frac{2k}{g}} \quad (4)$$

Prelab Questions

1. Derive Eq. (1) and write down an expression for the radius of gyration k in terms of the dimensions of the bar.
2.
 - a. Does Eq. (1) apply when:
 - i. A large amplitude is used?
 - ii. When damping is present, due to friction at the pivot, or to air resistance?
 - b. If Eq (1) does not apply, would the value you found for g be too high or too low?
3. Should the presence of holes in the bar be considered when calculating the theoretical value of k ?
4. Rewrite Eq (1) in the form $l^2 = f(T^2l)$ to give an equation of a straight line, with g related to the slope and k to an intercept.

5. Could there be a systematic error in the stopwatch you will use? How could you check this?
6. Will the increased weights used in the optional experiment alter the damping due to friction?

Apparatus

- pendulum bar and ball bearing mount
- 2 extra masses
- meter stick
- stopwatch

Experiment

1.
 - a. Determine the period of the compound pendulum for various values of l . To do this, time about 20 complete swings and repeat each measurement several times. Be careful not to make the amplitude of oscillation too large and explain why this precaution is necessary.
 - b. Plot your results and calculate k from the minimum period.
 - c. Show that $l_1 l_2 = k^2$ for fixed T .
2. Calculate a theoretical value of k from the bar's dimensions and compare it with your experimental result from item 1.
3. Using the theoretical value of k , plot the theoretically predicted variation of the period with l on the graph on which you displayed your experimental results. Compare and comment.
4. Replot your data exploiting the linear relationship derived in the Prelab Questions and extract k and g from the graph. Compare to calculated or previously measured values.

Optional Experiment

Attach two equal masses symmetrically at each end of the bar. Repeat the experiment and interpret your results.